

General Agriculture Application Information

For

BERRIES

Soil, Foliar and Seed Applications

Products:

The Bio-N-Liven Answer®

The Carbon Answer®

The Mineral Electrolyte Answer®

Note: Application rates are “general.”
They have been compiled over the previous 15 years
from field tests and long-term usage by growers.

Such tests are ongoing and this table will
be updated as more information
is made available.





THE ANSWER® 2005 PRODUCT APPLICATION RATES

BERRIES

Application Area	The Answer® Products	Application Rate	Application Time	Dilution Rate	Method of Application	Field Observations
Soil	Bio-N-Liven Answer®	Moderate Soil: (24 fl. Oz./acre)	On new crop before planting. In the spring	Ground - at least 1:100 Dilution rate may vary depending on spray equipment used.	Banded. Ground. Spray diluted product evenly in furrows or along bushes. After application, incorporate product into the soil. Rain or irrigation after application will improve performance.	INCREASED: Soil fertility, water holding capacity and drainage, soil aeration, neutralization of soil pH
Seed	Bio-N-Liven Answer®	For transplanted seedlings only (2 fl. Oz./acre)	Planting	At least 1:20. Use enough water, but not more than necessary to dip roots of plants to be planted on one acre. Calculate dilution rate so recommended amount will be used per acre.	Dip roots of seedlings in diluted product immediately after removing from seedbed and transplant in soil. Water transplanted seedlings with remaining solution.	IMPROVED: Survival of seedlings, regeneration and growth of root system, resistance of seedlings to stress (low temperature, drought, disease, etc.)
Seed	Mineral Electrolyte Answer®	For transplanted seedlings only (2 fl. Oz./acre)	Planting	At least 1:20. Use enough water, but not more than necessary to dip roots of plants to be planted on one acre. Calculate dilution rate so recommended amount will be used per acre.	Dip roots of seedlings in diluted product immediately after removing from seedbed and transplant in soil. Water transplanted seedlings with remaining solution.	IMPROVED: Survival of seedlings, regeneration and growth of root system, resistance of seedlings to stress (low temperature, drought, disease, etc.)

APPLICATION COLOR GUIDE

SOIL
 SEED
 FOLIAR



THE ANSWER® 2005 PRODUCT APPLICATION RATES

BERRIES

Application Area	The Answer® Products	Application Rate	Application Time	Dilution Rate	Method of Application	Field Observations
Foliar	Bio-N-Liven Answer®	(12 fl. Oz./acre)	Twice: At early bloom to 30% bloom and For multiple picking: After each harvest	Aerial - 1:20, 1:50 Ground - at least 1:100 Dilution rate may vary depending on the spray equipment used.	Spray diluted product evenly on plant foliage. For best uptake, apply product in evening and early morning. Heavy rain within 24 hours of application will diminish product effectiveness.	IMPROVED: Plant vigor, yield, quality of fruits
Foliar	Mineral Electrolyte Answer®	(3 fl. Oz./acre)	Twice: At early bloom to 30% bloom and For multiple picking: After each harvest	Aerial - 1:20, 1:50 Ground - at least 1:100 Dilution rate may vary depending on the spray equipment used.	Spray diluted product evenly on plant foliage. For best uptake, apply product in evening and early morning. Heavy rain within 24 hours of application will diminish product effectiveness.	IMPROVED: Plant vigor, yield, quality of fruits
Foliar	Carbon Answer®	(1-2 fl. Oz./acre)	Twice: At early bloom to 30% bloom and For multiple picking: After each harvest	Aerial - 1:20, 1:50 Ground - at least 1:100 Dilution rate may vary depending on the spray equipment used.	Spray diluted product evenly on plant foliage. For best uptake, apply product in evening and early morning. Heavy rain within 24 hours of application will diminish product effectiveness.	IMPROVED: Plant vigor, yield, quality of fruits

APPLICATION COLOR GUIDE



SOIL



SEED



FOLIAR